

University of M'sila

Faculty of: **Technology**

Common Base

First Series of exercises

Exercise 01:

Given the vectors \vec{V}_1 and \vec{V}_2 in an orthonormal basis $(\vec{i}, \vec{j}, \vec{k})$ such that:

$$\vec{V}_1 = \vec{i} + 2\vec{j} \quad \text{and} \quad \vec{V}_2 = 2\vec{i} - \vec{j}$$

1° Find the vector sum $\vec{S} = \vec{V}_1 + \vec{V}_2$, graphically and analytically.

2° Find the vector difference $\vec{D} = \vec{V}_1 - \vec{V}_2$ graphically and analytically.

3° The vectors \vec{V}_1 and \vec{V}_2 form a parallelogram. What represents graphically, the magnitude of the sum $|\vec{S}|$ and the magnitude of difference $|\vec{D}|$ in this parallelogram?

4° Determine the moduli of the vectors: $\vec{V}_1, \vec{V}_2, \vec{S}$ and \vec{D} .

Additional questions: If $\vec{A} + \vec{B} = 5\vec{i} - \vec{j}$ and $\vec{B} - \vec{A} = \vec{i} + \vec{j}$

5° Found the moduli of the vectors: $|\vec{A}|$, $|\vec{B}|$, $|\vec{A} + \vec{B}|$ and $|\vec{B} - \vec{A}|$?

6° Found the angles formed between: $(\vec{A}$ and $\vec{B})$; $(\vec{A} + \vec{B}$ and $\vec{A})$; $(\vec{B} - \vec{A}$ and $\vec{B})$;

$(\vec{A} + \vec{B}$ and $\vec{B} - \vec{A})$

7° Determine the components of \vec{n} the normal to the plane constituted by the vectors \vec{A} and \vec{B}

8° What are the components of \vec{A} and \vec{B} along the directions $\vec{u} = \vec{i} + \vec{j}$ and $\vec{v} = \vec{i} - \vec{j}$?

Exercise 02:

Given the vectors \vec{a} and \vec{b} in an orthonormal basis, $(\vec{i}, \vec{j}, \vec{k})$ such that:

$$\vec{a} = 3\vec{i} - 5\vec{j} + \vec{k} \quad \text{and} \quad \vec{b} = 2\vec{i} + 3\vec{j} - 4\vec{k}$$

1° Calculate the scalar (dot) product between \vec{a} and \vec{b} .

2° What is the angle between \vec{a} and \vec{b} . Determine $|\vec{a} + \vec{b}|$ and $|\vec{a} - \vec{b}|$ in two ways.

3° Determine the projection along the direction \vec{a} of the vector \vec{b}

If these vectors whose components are given according to the parameters α and β such that

$$\vec{a} = \alpha\vec{i} - 2\vec{j} + \vec{k} \quad \text{and} \quad \vec{b} = \beta\vec{i} + \vec{j} + \vec{k}$$

4°/ What is the relationship between α and β such that \vec{a} and \vec{b} are always perpendicular?

Exercise 03:

Given the vectors \vec{A} and \vec{B} in an orthonormal basis $(\vec{i}, \vec{j}, \vec{k})$

$$\vec{A} = 2\vec{i} - 3\vec{j} + 4\vec{k} \quad \text{and} \quad \vec{B} = \vec{i} + 5\vec{j} + 2\vec{k}$$

1°/ Calculate the vector (cross) product between \vec{A} and \vec{B} .

2°/ Find the angle between \vec{A} and \vec{B} .

3°/ What is the area constituted by the vectors \vec{A} and \vec{B} .

What is the direction of this surface?

If these vectors whose components are given according to the parameters γ and δ such that:

$$\vec{A} = \gamma\vec{i} - 3\vec{j} + 4\vec{k} \quad \text{and} \quad \vec{B} = 5\vec{i} + \delta\vec{j} + 2\vec{k}$$

4°/ What are the values of γ and δ so that \vec{A} and \vec{B} are always collinear?

Exercise 04:

In an orthonormal basis $(\vec{i}, \vec{j}, \vec{k})$, we give the vectors:

$$\vec{A}(t) = 2t\vec{i} + (t + 1)\vec{j} \quad \text{and} \quad \vec{B}(t) = 4t\vec{i} - 3t\vec{j} + 2\vec{k}$$

1°/ Calculate the derivatives $\frac{d\vec{A}}{dt}$, $\frac{d\vec{B}}{dt}$ of the vectors \vec{A} and \vec{B} .

2°/ Calculate derivatives $\frac{d(\vec{A} \cdot \vec{B})}{dt}$ and $\frac{d(\vec{A} \wedge \vec{B})}{dt}$ in two ways.

QCU:

1°/ Let be the vectors $\vec{A} = 3\vec{i} + 4\vec{j}$ and $\vec{B} = 7\vec{i} - 24\vec{j}$. The vector having the same modulus as \vec{B} and the same direction as \vec{A} is:

a/ $5\vec{i} + 20\vec{j}$ b/ $20\vec{i} + 15\vec{j}$ c/ $15\vec{i} + 10\vec{j}$ d/ $15\vec{i} + 20\vec{j}$

2°/ Let the vector $\vec{A} = 2\vec{i} + 3\vec{j}$. The angle between \vec{A} and the axis \vec{oy} is:

a/ $\arcsin \left[\frac{3}{2} \right]$ b/ $\arctg \left[\frac{3}{2} \right]$ c/ $\arctg \left[\frac{2}{3} \right]$ d/ $\arccos \left[\frac{3}{2} \right]$

3°/ 5 forces, each equal to '10N' and applied at the same point. These forces are coplanar and angles between each two consecutive forces are same. The resultant is:

a/ Zéro b/ 10N c/ 20N d/ $10\sqrt{2}N$

Exercise 05:

In an orthonormal basis (\vec{i}, \vec{j}) , we give the vector \vec{A} such that $\vec{A} = \vec{i} + \sqrt{3}\vec{j}$

1° Write the unit vector \vec{u}_A of \vec{A} in the base (\vec{i}, \vec{j})

This unit vector \vec{u}_A taken as a vector of the polar basis, $\vec{u}_A = \vec{u}_\rho$

2° Give the expression (in the Cartesian base) of the second vector of this base \vec{u}_θ .

3° Write the vector \vec{A} in the polar base.

Given a vector \vec{B} in the polar basis $\vec{B} = \rho\vec{u}_\rho + \sin\theta\vec{u}_\theta$

4° Give the expression of \vec{B} in the Cartesian base

Exercise 06:

Given a vector $\vec{A} = \vec{i} - \sqrt{3}\vec{j} - 2\vec{k}$

1° Give the spherical coordinates of \vec{A} ?

2° What is the spherical base $(\vec{u}_r, \vec{u}_\theta, \vec{u}_\phi)$ for \vec{A} , expressed in the Cartesian basis?

3° Do the same thing again for the vector \vec{A} in the cylindrical base $(\vec{u}_\rho, \vec{u}_\theta, \vec{k})$.

Exercise 07:

In an orthonormal basis (\vec{i}, \vec{j}) , we give the point $M\left(\begin{smallmatrix} \sqrt{3} \\ 1 \end{smallmatrix}\right)$ on the circle of radius $R = 2$ and center $C(0, 0)$:

1° Write the unit vectors of the polar basis $(\vec{u}_\rho, \vec{u}_\theta)$ in the Cartesian basis (\vec{i}, \vec{j}) .

Given \vec{u}_ρ and \vec{u}_θ for the point $M\left(\begin{smallmatrix} \sqrt{3} \\ 1 \end{smallmatrix}\right)$.

2° Write the derivatives $\frac{d\vec{u}_\rho}{dt}$ and $\frac{d\vec{u}_\theta}{dt}$ of the unit vectors $\vec{u}_\rho, \vec{u}_\theta$ in the same polar basis if

$$\frac{d\theta}{dt} = \dot{\theta} = t \quad .$$

3° Write the unit vectors of the intrinsic (\vec{u}_T, \vec{u}_N) basis in the Cartesian basis (\vec{i}, \vec{j}) .

Given \vec{u}_T and \vec{u}_N for the point $M\left(\begin{smallmatrix} \sqrt{3} \\ 1 \end{smallmatrix}\right)$.

4° Write derivatives $\frac{d\vec{u}_T}{dt}$ and $\frac{d\vec{u}_N}{dt}$ of the unit vectors in the same intrinsic basis \vec{u}_T, \vec{u}_N .

5° Represent the polar and intrinsic basis at the point $M\left(\begin{smallmatrix} \sqrt{3} \\ 1 \end{smallmatrix}\right)$

Exercise 08: (Additional)

Let a vector $\vec{A} = 3\vec{i} + 2\vec{j} + \vec{k}$

1°/ Give the spherical coordinates of \vec{A} ?

2°/ Write the expressions of the spherical base $(\vec{u}_r, \vec{u}_\theta, \vec{u}_\varphi)$, in the Cartesian base

3°/ Do the same thing again for the vector \vec{A} in the cylindrical base $(\vec{u}_\rho, \vec{u}_\theta, \vec{k})$.

Exercise 09: (HW)

1°/ Express the Cartesian base $(\vec{i}, \vec{j}, \vec{k})$ in the spherical base $(\vec{u}_r, \vec{u}_\theta, \vec{u}_\varphi)$

2°/ Show that the unit vectors of the spherical basis are written as follows:

$$\frac{d\vec{u}_r}{dt} = \vec{\Omega}_1 \wedge \vec{u}_r \quad \frac{d\vec{u}_\theta}{dt} = \vec{\Omega}_2 \wedge \vec{u}_\theta \quad \frac{d\vec{u}_\varphi}{dt} = \vec{\Omega}_3 \wedge \vec{u}_\varphi.$$

Give the expression of $\vec{\Omega}_1$, $\vec{\Omega}_2$ and $\vec{\Omega}_3$.