

Mohamed Boudiaf University - M'sila
 Faculty of Technology
 Department of Civil Engineering- Department of
 Electrical Engineering
 Module: Probability-Statistics
**Chapter 3 : COMBINATORIAL
 ANALYSIS**

Merini Abdelaziz*

19 novembre 2024

Table des matières

1	Introduction	2
2	Preliminaries	2
2.1	Definitions :	2
2.2	Factor notation	2
3	Basic Counting Principles	2
3.1	Addition Principle	3
3.2	Multiplication Principle	3
4	Permutations	4
4.1	Permutations of n distinct elements	4
4.2	Permutations with Similar Elements	4
4.3	Circular Permutations	5
4.4	r-Permutation (arrangement)	5
5	Combinations	6
5.1	Combination	6
5.2	Combinations with Repetition	7

*

5.3	Pascal's Trianlge	7
5.4	Newton's Binomial Theorem	8

1 Introduction

Combinatorial analysis is the science of counting and is used to determine the number of possible outcomes of a given experiment. Knowledge of these enumeration methods is essential for the elementary calculation of probabilities.

2 Preliminaries

2.1 Definitions :

- A set Ω is finite when it has a finite number of elements. The number of elements of Ω is called the cardinal of the set and it is noted : $\#\Omega$ or $|\Omega|$.
- To count is to determine the number of elements in a finite set, i.e. to determine its cardinal.

Example 1. If $\Omega = \{heads, tails\}$, its cardinal is $|\Omega| = 2$.
 - If $\Omega = \{1, 2, 3, 4, 5, 6\}$, $|\Omega| = 6$.
 If Ω is the set of integers, $|\Omega| = +\infty$. $card(\emptyset) = 0$.
 The set Omega of players in a football team is a finite set. . Then $|\Omega| = 11$.

Definition 1. We say that two sets Ω_1 and Ω_2 are **disjoint**, if they have no elements in common, i.e.

$$\Omega_1 \cap \Omega_2 = \emptyset$$

2.2 Factor notation

Definition 2. Let $n \in \mathbb{N}$. The product of the integers from 1 to n is called n factorial, denoted $n!$

$$n! = \begin{cases} 1 & \text{if } n = 0 \\ n \times (n - 1) \times (n - 2) \times \dots \times 3 \times 2 \times 1 & \text{if } n > 0 \end{cases}$$

Example 2. $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$. $6! = 6 \times 5! = 720$.
 $0! = 1$.

Rremark 1. $\frac{n!}{(n - 1)!} = n$, $n! = n \times (n - 1)!$

3 Basic Counting Principles

The basic counting principles are **the product rule** and **the sum rule** will be presented and show how can be used to solve many different counting problems.

3.1 Addition Principle

Suppose that a set Ω is partitioned into pairwise disjoint parts $\Omega_1, \Omega_2, \dots, \Omega_k$

The addition principle claims that in this case

$$|\Omega| = |(\Omega_1 \cup \Omega_2 \cup \dots \cup \Omega_k)| = |(\Omega_1)| + |(\Omega_2)| + \dots + |(\Omega_k)|$$

Example 3. Let $\Omega_1 = \{a, b, c, d\}$ and $\Omega_2 = \{g, h, k\}$. we have $\Omega_1 \cap \Omega_2 = \phi$, then

$$|(\Omega_1 \cup \Omega_2)| = |(\Omega_1)| + |(\Omega_2)| = 4 + 3 = 7.$$

Example 4. Let Ω be the set of students attending the combinatorics lecture. It can be partitioned into parts Ω_1 and Ω_2 where :

Ω_1 = set of students that like easy examples.

Ω_2 = set of students that don't like easy examples.

If $|\Omega_1| = 22$ and $|\Omega_2| = 8$ then we can conclude $|\Omega| = 30$.

3.2 Multiplication Principle

If k experiments that are to be performed are such that :

- the first one may result in any of n_1 possible outcomes ;
- and if, for each of these n_1 possible outcomes, there are n_2 possible outcomes of the second experiment ;
- and if, for each of the possible outcomes of the first two experiments, there are n_3 possible outcomes of the third experiment ;
- and if ...,

then there is a total of $n_1 \times n_2 \times \dots \times n_k$ possible outcomes of the k experiments.

Example 5. We want to print a car number plate with, from left to right, 2 distinct letters and 3 digits, the first of which is not zero (a global experiment). How many plates of this type are there ? (CH124, DE665,,) .

Solution 1.

<i>letter</i>	<i>letter</i>	<i>number</i>	<i>number</i>	<i>number</i>
↑	↑	↑	↑	↑
26	25	9	10	10

According to the Multiplication Principle, the possible number of plates of this type is

$$26 \times 25 \times 9 \times 10 \times 10 = 585000$$

Example 6. Three 6-sided dice are rolled in succession (a global experiment). How many possible outcomes are there ?

Solution 2. For D_1 , we have (6 distinct numbers)

For D_2 , we have (6 distinct numbers)

For D_3 , we have (6 distinct numbers)

According to the Multiplication Principle, the number of possible results is

$$6 \times 6 \times 6 = 216.$$

4 Permutations

4.1 Permutations of n distinct elements

Definition 3. A permutation of n distinct elements is an ordered arrangement where each element is used once.

Example 7. The possible permutations of the 3 letters a, b, c are :
 $abc, bca, cab, bac, acb, cba$.

Proposition 1. The number of permutations with n distinct elements is

$$P(n, n) = n!$$

Rremark 2. Use **permutations** : if a problem calls for the number of arrangements of objects and different orders are to be counted.

Example 8. In the example 7, on a $n = 3$, donc $P_3 = 3! = 6$.

4.2 Permutations with Similar Elements

Definition 4. If we classify in a particular order n elements of which
 n_1 are identical of type 1,
 n_2 are identical of type 2,
.....,
 n_k are identical of type k ,
we form a permutation with repetitions of these $n = n_1 + n_2 + \dots + n_k$ elements.

Proposition 2. The number of permutations with repetitions is

$$\bar{P}_n(n_1, n_2, \dots, n_k) = \frac{n!}{n_1!n_2!\dots n_k!}$$

Example 9. How many different words can you make with all the letters in the word « ERRER »

We have 5 letters ($n = 5$) some of which are similar.

the letter 'E' appears twice so $n_1 = 2$, the letter 'R' appears three times so $n_2 = 3$, then

$$\bar{P}_5(2; 3) = \frac{5!}{2!3!} = 10$$

RRREE, RRERE, RREER, RERRE, RERER, REERR, ERRRE, ERRER;
ERERR; EERRR.

Example 10. A chess tournament has 10 competitors, of which 4 are Russian, 3 are from the United States, 2 are from Great Britain, and 1 is from Brazil. If the tournament result lists just the nationalities of the players in the order in which they placed, how many outcomes are possible?

Solution There are $\frac{10!}{2!3!1!1!1!1!} = 12600$ possible outcomes

Rremark 3. a) $\bar{P}_n(n_1, n_2, \dots, n_k) < P_n$.

b) The bar above the P stands for 'with repetition'.

4.3 Circular Permutations

Definition 5. Circular permutation is the total number of ways in which n distinct objects can be arranged around a fixed circle.

Proposition 3. The number of circular permutations of n objects is equal to

$$(n - 1)!$$

Example 11. How many ways can you arrange 4 people :

a) On a line ?

b) Around a round table ?

Solution 3. a) $5! = 120$.

b) $(5-1)! = 4! = 24$.

4.4 r-Permutation (arrangement)

Definition 6. Let r be a positive integer ($r \leq n$). By an r -permutation of a set Ω of n elements, we understand an ordered arrangement of r of the n elements.

Example 12. If $\Omega = \{a, b, c\}$ then the three 1-permutations of Ω are a, b, c , the six 2-permutations of S are ab, ac, ba, bc, ca, cb , and the six 3-permutations of S are $abc, acb, bac, bca, cab, cba$.

Proposition 4. The number of r -permutations from a set containing n elements is given by

$$\begin{aligned} P(n, r) &= \frac{n!}{(n-r)!} \\ &= n(n-1) \dots (n-r+1) \end{aligned}$$

Rremark 4. If $r = n$, we have $P(n, r) = \frac{n!}{(n-n)!} = \frac{n!}{0!} = n!$,

Example 13. With the letters of the word « RELATION », how many can you form, with or without meaning, different 5-letter words ?

Answer : $n = 8$ et $r = 5$, there are $A_8^5 = \frac{8!}{(8-5)!} = 8 \times 7 \times 6 = 336$ possible words

Proposition 5. *The number of permutations of n objects, taken r at a time, when repetition of objects is allowed, is*

$$n^r$$

Example 14. *How many 3 digit numbers can be formed with the digits 1 and 2?*

Answer : $n = 2$, there are $2^3 = 2^3 = 8$. numbers
111, 112, 121, 211, 122, 222, 221, 212.

Example 15. *How many two-digit numbers can you make using the digits 5, 6, 7, 8, 9?*

Answer : $n = 5$, $r = 2$, $5^2 = 25$.

5 Combinations

5.1 Combination

Definition 7. *A combination of a set of elements is an arrangement where each element is used once, and order is not important.*

Example 16. *Four people {1; 2; 3; 4} want to play doubles table tennis. How many different teams can they form?*

Answer : *It's a combination : {1; 2} , {3; } , {1; 3} , {2; 4} , {1; 4} , {2 : 3} We can form 6 teams.*

Proposition 6. *The Number of Combinations of n Objects Taken r at a Time*

$$\begin{aligned} \binom{n}{r} &= \frac{n!}{r!(n-r)!} \\ &= \frac{P(n, r)}{r!} \end{aligned}$$

Rremark 5. 1) *We also note $\binom{n}{r}$.*

2) *Use combinations if a problem calls for the number of ways of selecting objects and the order of selection is not to be counted.*

Example 17. *A committee of 3 is to be formed from a group of 24 people. How many different committees are possible?*

Solution

There are

$$\binom{24}{3} = \frac{24!}{3!21!} = \frac{24 \times 23 \times 22}{6} = 2024$$

possible committees

Properties

- 1) $\binom{n}{n} = \binom{n}{0}$, $C(n, r) = C(n, n - r)$ (symmetry formula)
- 2) $\binom{n}{1} = \binom{n}{n-1} = n$.
- 3) $\binom{n+1}{r+1} = \binom{n}{r} + \binom{n}{r+1}$ (Pascal's triangle).
- 4) $\sum_{i=0}^n \binom{n}{i} = 2^n$

Démonstration. exercice. □

Example 18. $\binom{3}{0} = \frac{3!}{0!(3-0)!} = 1$; $\binom{3}{3} = 1$; $\binom{3}{2} = \binom{3}{1} = 3$

5.2 Combinations with Repetition

Definition 8. A combination with repetition of r objects from n is a way of selecting r objects from a list of n . The selection rules are :

- 1) the order of selection does not matter
- 2) each object can be selected more than once.

Example 19. Let $\Omega = a, b, c$. We want to obtain the list of combinations of $k=4$ elements taken from the set E , i.e. generate the list of 4-combinations with repetition :

(a, a, a, a) , (a, a, a, b) , (a, a, a, c) , (a, a, b, b) , (a, a, b, c) , (a, a, c, c) ,
 (a, b, b, b) , (a, b, b, c) , (a, b, c, c) , (a, c, c, c) , (b, b, b, b) , (b, b, b, c) ,
 (b, b, c, c) , (b, c, c, c) , (c, c, c, c)

Proposition 7.

Number of combinations with repetition of r objects from n is

$$\binom{n+r-1}{r} = \frac{(n+r-1)!}{r!(n-1)!}$$

Example 20. In the previous example , we have $\binom{3+4-1}{4} = \binom{6}{4} = 15$

5.3 Pascal's Triangle

The relationship $\binom{n+1}{r+1} = \binom{n}{r} + \binom{n}{r+1}$

allows practical determination of the various coefficients $\binom{n}{r}$ using Pascal's triangle.

If $r > n$, , we put $\binom{n+r-1}{r} = 0$

$n \setminus r$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

5.4 Newton's Binomial Theorem

Let n be a positive integer. Then, for all x and y ,

$$(a + b)^n = \sum_{i=0}^n \binom{n}{i} a^{n-i} b^i$$

Example 21. $(a + b)^2 = \binom{2}{0} a^2 + \binom{2}{1} ab + \binom{2}{2} a^0 b^2 = 1a^2 + 2ab + 1b^2$

$$\begin{aligned} (a + b)^3 &= \binom{3}{0} a^3 + \binom{3}{1} a^2 b + \binom{3}{2} a^1 b^2 + \binom{3}{3} a^0 b^3 \\ &= 1a^3 + 3a^2 b + 3a^1 b^2 + 1b^3. \end{aligned}$$